A new species of the genus *Pseudicius* Simon 1885 (Araneae: Salticidae) from SW Iran

Dmitri V. Logunov

The Manchester Museum, The University of Manchester, Oxford Road, Manchester M13 9PL, UK

E-mail: dmitri.v.logunov@manchester.ac.uk

Abstract — A new species *Pseudicius admirandus* sp. nov. (\mathscr{E}°) is described from the Khūzestān Province, SW Iran.

Key words — Spiders, description, Khūzestān

Introduction

The Salticidae fauna of Iran is known to number 70 species of 35 genera (Logunov *et al.* 2001; Sahra 2006). In the present paper a new species of *Pseudicius* collected from Khūzestān Province, one of the least studied regions of Iran, is described. The terminology and format of the description follows Logunov (2001). The type specimens are kept in the Zoological Museum of the Moscow State University, Moscow, Russia (ZMUM; Dr K.G. Mikhailov). Comparative type material on *P. rudakii* Prószyński 1992 was borrowed from the National Museum of Natural History, Smithsonian Institution, Washington, USA (NMNH; Dr. J. Coddington and Ms D. de Rocha).

Description

Pseudicius admirandus sp. nov. (Figs 1–5)

Type material. The ♂ holotype and ♀ paratype (ZMUM) from Iran, Khūzestān, Ahwaz (=Ahvāz; c. 31°19′N, 48°4 1′E), Sheyban, 7.7.2005, B. Vazirianzadeh, Samoravej & Shirsian.

Comparative material. The & holotype of Pseudicius rudakii Prószyński 1992 (NMNH; Figs. 6, 7) from Iran, Fars, Shiraz, Park Hotel, 1.05.1965, I. Neal.

Etymology. The specific epithet derives from the Latin word 'admirandus', meaning 'wonderful, wondrous'.


Diagnosis. P. admirandus sp. nov. belongs to the cinctus species group (sensu Prószyński 2007). Its male is most similar to P. rudakii described and known after a single male from Iran (Prószyński 1992; Figs. 6, 7), from which it can be easily distinguished by the wider and stronger embolus (cf. Figs 1 and 6) and the shorter and narrower verntral branch of the tibial apophysis (cf. Figs. 2 and 7). The female copulatory organs of P. admirandus sp. nov. are similar to those of P. ludhianaensis (Tikader 1974) described from Punjab, India (see Tikader 1974: figs. 1–3, sub.

Marpissa l.; and Sadana, 1980: figs. 1–5, sub. Phlegra citri). Epigynes of both species have the long median septum, whereas the spermathecae of *P. ludhianaensis* was not illustrated. Despite this, females of these species can be separated by the dorsum pattern (three longitudinal orange stripes in *P. admirandus* sp. nov. and two transverse brown bands in *P. ludhianaensis*; Tikader 1974: fig. 1 and Sadana, 1980: fig. 1).

Distribution. The type locality only.

Description. MALE (holotype). Measurements. Carapace 2.25 long, 1.50 wide, 0.75 high at PLE. Ocular area 0.98 long, 1.13 wide anteriorly and 1.23 wide posteriorly. Diameter of AME 0.38. Abdomen 2.68 long, 1.38 wide. Chelicera length 0.75. Clypeus height 0.10. Length of leg segments: I 1.28 + 0.88 + 1.25 + 0.68 + 0.38; II 0.98 + 0.58 + 0.65 + 0.53 + 0.30; III 1.00 + 0.55 + 0.60 +0.70 + 0.40; IV 1.15 + 0.63 + 0.86 + 0.85 + 0.40. spination. Leg I: Fm d 1-1-1, pr 2ap; Tb pr 0-0-1, v 0-1-1-1ap; Mt v 2-2ap. Leg II: Fm d 1-1-1, pr 2ap; Tb pr 0-0-1, v 1ap; Mt v 2-2ap. Leg III: Fm d 1-1-2, pr 0-0-1-2; Pt rt 0-1-0; Tb pr 0-1, rt 1-1, v 2ap; Mt pr and rt 2ap, v 0-1-2ap. Leg IV: Fm d 1-1-3; Tb pr 0-0-1, rt 1-1, v 1-0-2ap; Mt pr and rt 2ap, v 1-0-2ap. Coloration. Carapace yellow, with two longitudinal wide orange stripes along the ALE-PLE line. Clypeus narrow, covered with orange scales. Black around eyes. The row of stridulatory bristles beneath eyes present on each side of the carapace. Sternum, maxillae, labium yellow. Abdomen gray-yellow; dorsum with one medial and two lateral longitudinal wide orange stripes; venter yellow, with gray reticulate pattern. Book-lung covers yellow, spinnerets gray brownish. All legs and palps yellow, but Pt I anterior-distally and Tb I anterior- and porterior-distally brownish. Palpal structure as in Figs. 1, 2.

FEMALE (paratype). *Measurements*. Carapace 2.05 long, 1.43 wide, 0.60 high at PLE. Ocular area 0.88 long, 1.20 wide anteriorly and 1.20 wide posteriorly. Diameter of AME 0.35. Abdomen 3.18 long, 1.45 wide. Chelicera length 0.53. Clypeus height 0.10. Length of leg segments:



Figs. 1–7. Copulatory organs of *Pseudicius admirandus* sp. nov., the 3 holotype and 9 paratype (1–5) and *P. rudakii* Prószyński, 1992, the 3 holotype (6, 7). — 1, 6, male palp, verntral view; 2, 7, ditto, retrolateral view; 3, epigyne, ventral view; 4, spermathecae, dorsal view; 5, ditto, ventral view. Scales: 0.1 mm.

I 0.88 + 0.60 + 0.65 + 0.40 + 0.35; II 0.78 + 0.50 + 0.48 + 0.40 + 0.38; III 0.93 + 0.50 + 0.55 + 0.60 + 0.40; IV 1.15 + 0.53 + 0.75 + 0.68 + 0.53. Leg spination. Leg I: Fm d 0-1-1-2; Tb pr 0-0-1-1ap, v 0-1-1-1ap; Mt v 2-2-2ap. Leg III: Fm d 0-1-1-2; Tb v 0-0-1-1ap; Mt v 2-2-2ap. Leg III: Fm d 0-1-1-2; Tb pr and rt 0-0-1, v 2ap; Mt pr 2ap, rt and v 0-1-2ap. Leg IV: Fm d 0-1-1-3; Tb pr and rt 1ap, v 1-2ap; Mt pr, rt and v 2ap. Coloration. Carapace as in males, except as follows: clypeus covered with white scales; dorsum without longitudinal orange stripes, but instead with an orange reticulate pattern. Epigyne and spermathecae as in Figs. 3-5.

Acknowledgments

I wish to express my warmest thanks to Dr J. Coddington and Ms Dana de Roche (of the NMNH) for giving access to the spider collections under their care.

References

Logunov, D. V. 2001. New and poorly known species of the jumping spiders (Aranei: Salticidae) from Afghanistan, Iran and Crete. Arthropoda Selecta 10(1): 59-66.

Logunov, D. V., Marusik Yu.M. & Mozaffarian F. 2001. Faunistic review of the jumping spiders of Iran (Aranei: Salticidae). Arthropoda Selecta, 10(2): 155–167.

Prószyński, J. 1992. Salticidae (Araneae) of the Old World and

- Pacific Islands in several US collections. Annales zoologici, Polska Akademia Nauk, 44(8): 87-163.
- Prószyński, J. 2007. Monograph of the Salticidae (Araneae) of the World; the page on *Pseudicius* last updated Feb 2006. Online at: http://www.miiz.waw.pl/salticid/main.htm
- Sahra, G. 2006. Renewed Checklist of spiders (Aranei) of Iran. Pakistan Journal of Biological Sciences, 9(10): 1839–1851.
- Sadana, G. L. 1980. A new Indian species of Phlegra Simon
- (Araneida: Salticidae). Entomologist's monthly Magazine, 115: 229-230.
- Tikader, B. K. 1974. Studies on some jumping spiders of the genus *Marpissa* from India (family-Salticidae). Proceedings of the Indian Academy of Sciences, 79(5, Sec.B): 204–215.

Received April 4, 2007 / Accepted April 28, 2007